

53883-172

11/15/2012

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Jane Miller
Control Solutions, Inc
c/o Biologic, Inc
115 Obtuse Hill Road
Brookfield, CT 06804

NOV 15 2012

Subject: Label amendment adding optional marketing statements and correct use rates.
Prodiamine 0.37 Plus
EPA Reg. No: 53883-172
Application Dated: August 16, 2012

Dear Ms. Miller

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

Please note that marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available, then those claims must be removed.

A stamped copy of your label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Grant Rowland at (703) 347-0254 or at Rowland.Grant@epa.gov.

Sincerely,

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

08 06 2012 Label amendment to add marketing statements and correct application rate.

PRODIAMINE 0.37 PLUS

[with Lawn Fertilizer]

Optional Marketing Language:

[FOR SELECTIVE PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS]

[STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL & MORE]

[Prevents Crabgrass in Lawns]

[Phosphorus-Free][Helping our water sources]

[With Lawn Fertilizer]

[Controls crabgrass and broadleaf weeds before they germinate]

[Feeds for 3 Months guaranteed!]

[Prevent crabgrass and fertilize your lawn in one application]

[Up to 5 month weed and grass control]

[Covers 5,000 sq. ft.] for 15 lb. bag

ACTIVE INGREDIENT:

Prodiamine: [N³,N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]

0.37%

INERT INGREDIENTS:

99.63%

TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-172

EPA EST. NO.

Control Solutions, Inc.
5903 Genoa-Red Bluff Rd.
Pasadena, TX 77507-1041
281-892-2500

NET CONTENT: _____

RECEIVED
NOV 15 2012
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
this pesticide is registered.
53883-172

08 06 2012 Label amendment to add marketing statements and correct application rate.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> ● Call a poison control center or physician immediately for treatment advice. ● Have person sip a glass of water if able to swallow. ● Do not induce vomiting unless told to do so by the poison control center or physician. ● Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> ● Move person to fresh air. ● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. ● Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.</p>	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

08 06 2012 Label amendment to add marketing statements and correct application rate.

This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹In many areas a single application of 3.2 to 9.3 pounds/1,000 square feet (144 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 3.2 to 6.2 pounds/1,000 square feet (144 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).


²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

CRABGRASS SEED GERMINATION DATES

Approximate Date



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural

08 06 2012 Label amendment to add marketing statements and correct application rate.

practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate¹
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-Type)	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	6.2 lbs/1,000 sq. ft. per year (270 lbs/acre per year)
Creeping Red Fescue	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.6 lbs/1,000 sq. ft. per year (202 lbs/acre per year)
Creeping Bentgrass	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	4.0 lbs/1,000 sq. ft. per year (175 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	5
4.9 pounds/1,000 sq. ft. (215 pounds/acre)	-	6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)	-	-	9
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at

08 06 2012 Label amendment to add marketing statements and correct application rate.

the rate of 2.7 to 9.3 pounds/1,000 square feet (118 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.
Spreader A	X	X	X	X	X	X
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

COMMON NAME

- Abies spp.
- Abelia grandiflora
- Acer palmatum
- A. Platanoides
- Achillea spp.
- Actinidia chinensis
- Agapanthus africanus
- Agapanthus orientalis
- Akebia quintata
- Allium cernuum
- Anemone hybrida
- Aquilegia spp.
- Arctostaphylos densiflora
- Arctotheca calendula
- Artemisia spp.
- Aster spp.
- Aster X Frikartii
- Athrium Filiz-femina
- Aucuba japonica
- Begonia spp.
- Berberis gladwynensis
- B. Julianae
- B. mentorensis
- B. Thunbergii
- B. Verruculosa

- Fir species** (Balsam, Frasier, Noble, etc.)
- Abelia: Sherwood
- Japanese Maple
- Norway Maple
- Yarrow: King Edward
- Kiwi*
- Lily-of-the-Nile (African Lily)
- Five-leaf or Chocolate Vine
- Lady's Leek, Nodding Onion
- Japanese Anemone
- Aquilegia: Red and Gold
- Vine Hill Manzanita
- Cape Weed
- Wormwood; Silver Mound, Castle
- Aster: Bonnie Blue, Purple Dome
- Lady Fern; Fern Lady
- Japanese Aucuba
- Fibrous Begonia: Hardy Grandis
- Barberry
- Wintergreen Barberry
- Mentor Barberry
- Japanese Barberry
- Warty Barberry

08 06 2012 Label amendment to add marketing statements and correct application rate.

- Bergenia cordifolia
- Boltonia asteroides
- Buddleia davidii
- Buxus microphylla
- Callistemon viminalis
- Calluna vulgaris
- Campanula carpatica
- Campis X Tagliabuana
- Carpobrotus edulis
- Cassia artemisoides
- Ceanothus rigidus
- Ceratostigma plumbagonoides
- Chamaecyparis pisifera
- Cleyera japonica
- Citrus spp.
- Coreopsis spp.
- Cornus florida
- C. Stolonifera
- Cortaderia selloana
- Cotoneaster apiculatus
- C. buxifolius
- C. dammeri
- C. Microphyllus
- Cretaeagus spp.
- Crocsmia spp.
- Cupressus sempervirens
- Delosperma alba
- Delosperma spp.
- Delphinium spp.
- Dianthus deltoides
- D. gratianopolitanus
- Dodonea viscosa
- Echinacea purpurea
- Elaeagnus pungens
- Euonymus fortunei
- E. japonica
- E. kiautschovica
- Fatsia japonica
- Forsythia intermedia
- Forsythia suspensa
- F. viridissima
- Gaillardia spp.
- Gardenia jasminoides
- Gaura spp.
- Gentiana dahurica
- Geranium cinereum
- Gladiolus spp.
- Gypsophilia repens
- Hedera helix
- Hellanthemum spp.
- Hemerocallis spp.
- Hibiscus
- Hibiscus spp.
- Hibiscus rosa-sinensis
- Hosta plantaginea
- H. Sieboldiana
- Snowbank
- Butterfly-Bush (Dwarf Blue); Royal Red
- Japanese Boxwood
- Weeping Bottlebrush
- Scotch Heather
- Tussock Bellflower, (White Clips)
- Trumpet Flower, Madame Galen
- Hottentot Fig (Ice Plant)
- Feathery Cassia
- Wild Lilac
- False Cypress
- Cleyera
- Citrus species*
- Coreopsis (Calliopsis) Early Sunrise, Moonbeam
- Flowering Dogwood
- American Dogwood
- Pampas Grass
- Cranberry Cotoneaster
- Cotoneaster
- Bearberry Cotoneaster
- Rockspray Cotoneaster
- Hawthorne
- Lucifer
- Italian Cypress
- White Trailing Ice Plant
- Cooperi Pink
- Larkspur: Blue Elf
- Dianthus: Maiden Pinks "Zing"
- Cheddar Pink
- Hop Bush
- Coneflower, Purple; Magnus
- Silverberry
- Wintercreeper
- Evergreen Euonymus
- Spreading Euonymus
- Japanese Aralia
- Border Forsythia
- Weeping Forsythia
- Greenstem Forsythia
- Gaillardia, Blanket Flower: "Goblin"
- Gardenia, Cape-Jasmine
- Gentian
- Cranesbill
- Gladiolus species**
- Baby's Breath
- English Ivy
- Sunrose
- Daylily: Aztec Gold, Stella de Oro, Tender Love
- Rose of Sharon**
- Mallow: Disco Belle White
- Chinese Hibiscus
- Hosta, Plantain Lily (Fragrant)
- Hosta, "Seersucker"

08 06 2012 Label amendment to add marketing statements and correct application rate.

Houttuynia cordata var. variegata	Bigleaf Hydrangea
Hydrangea macrophylla	Holly
Ilex Bernyi	Chinese Holly
Ilex cornuta	Japanese Holly
I. crenata	American Holly
I. opaca	Yaupon Holly, Schillings
I. vomitoria	
Inula ensifolia	Sword-Leaved Iris; Jodlesong
Iris ensata	Siberian Iris; Cabernet
Iris-siberica	Iris species**
Iris spp.	Winter Jasmine
Jasminium nudiflorum	Chinese Juniper; Nick's Compact, Parsonnii
Juniperus chinensis	Shore Juniper; Blue Pacific
J. conferta	Parsoni
J. davurica	Creeping Juniper
J. horizontalis	Walnut*
Juglans spp.	Shrimp Plant
Justicia brandegeana	Crape Myrtle
Lagerstromia indica	Crape Myrtle; Tuscarora
Lagerstromia indica X Fauriei	Weeping Lantana
Lantana montevidensis	Lavender; Munstead
Lavender spp.	Edelweiss
Leonopodium alpinum	Amur Privet
Ligustrum amurense	Chinese Privet; Variegata
Ligustrum sinense	Japanese Privet
L. japonicum	Glossy Privet (Waxleaf)
L. lucidum	Lily; Jazz
Lilium spp.	Big Blue Lily, Silver Mound, Evergreen Giant
Liriope muscari	Liriope, Variegated
Liriope muscari var. variegata	Liriope, Creeping
Liriope spicata	Cardinal Flower, Indian Pink
Lobelia Cardinalis	Japanese Honeysuckle
Lonicera japonica	Tatarian Honeysuckle
L. tatarica	Burgundy
Lorapetalum chinense	Loosestrife; Modern Pink
Lythrum spp.	Magnolia species**
Magnolia spp.	Ice Plant
Maleophora luteola	Crabapple*
Malus spp.	Yaku Jima, Silberfeder
Miscanthus sinensis	Heavenly Bamboo
Nandina domestica	Narcissus species**
Narcissus spp.	Oleander
Nerium oleander	Olive*
Olea europaea	Mondo Grass
Ophiopogon japonicus	Osmanthus (False Holly); Gulf Tide
Omanthus heterphyllus	Trailing African Daisy
Osteospermum fruticosum	Sourwood
Oxydendrum arboretum	Tree Peony
Paeonia suffruticosa	Fountain Grass (Dwarf)
Pennisetum setaceum	
Perovskia atriplicifolia	Avocado*
Persea americana	Frasier's Photinia (Redtip)
Photinia Fraseri	Dragonhead, False; Vivid
Physotegia Virginiana	Spruce species** (Colorado Blue, Norway, etc.)
Picea spp.	Lily-of-the-Valley Bush
Pieris japonica	

08 06 2012 Label amendment to add marketing statements and correct application rate.

- Pinus brutia
- P. canariensis
- P. Elliottii
- P. halepensis
- P. nigra
- P. palustris
- P. radiata
- P. strobus
- P. sylvestris
- P. taeda
- P. Thunbergiana
- P. Virginiana
- Pistachio spp.
- Pittosporum rhombifolium
- P. tobira
- Podocarpus macrophyllus
- Prunus laurocerasus
- Prunus spp.

- Pseudotsuga menziesii
- Pyracantha coccinea
- P. fortuneana
- P. koidzumii
- Pyrus spp.
- Quercus Shumardii
- Quercus spp.
- Raphiolepis indica
- Raphiolepis umbellata
- Rhododendron
(including Azalea)

- Rosa banksiae
- Rosmarinus officinalis
- Rudbeckia spp.
- Santolina virens
- Saxifraga spp.
- Scabiosa spp.
- Sedum album
- Sedum caudicola
- Sedum dasyphyllum
- Sedum spurium
- Spiraea bumalda
- Syzygium paniculatum
- Taxus cuspidata
- T. media
- Teucrium spp.
- Thalictrum dipterocarpum
- Thuja occidentalis
- Trachelospermum asiaticum
- Tsuga canadensis
- Tulipa spp.
- Veronica spp.
- Viburnum japonicum
- V. odoratissimum
- V. plicatum

- Calabrian Pine
- Canary Island Pine
- Slash Pine
- Aleppo Pine
- Austrian Pine
- Longleaf Pine
- Monterey Pine
- Eastern White Pine
- Scotch Pine
- Loblolly Pine
- Japanese Black Pine
- Virginia Pine
- Pistachio*
- Queensland Pittosporum
- Japanese Pittosporum
- Japanese Yew
- English Laurel
- Almond, Apricot, Nectarine, Peach, Plum and Prune*
- Douglas Fir**
- Firethorn (Scarlet)
- Firethorn
- Firethorn
- Bradford Pear spp.
- Oak, Shumard's Red
- Oak species
- Indian Hawthorne
- Yedda Hawthorne
- 'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'
- 'Delaware Valley White,' 'Roseum Elegans,'
- 'Flame Creeper', 'Girard Crimson', 'Wakeiebisu'
- 'White Gumbo', 'George L. Tabor'
- Lady Bank's Rose
- Rosemary*
- Black-Eyed Susan; Goldstum
- Santolina
- Saxifrage; Purple Dome
- Pincushion Flower
- Stonecrop
- Stonecrop; Lidakense
- Stonecrop
- Stonecrop; Dragon's Blood
- Spirea; Anthony Waterer
- Australian Brushcherry
- Japanese Yew
- Yew
- Germander
- Meadow Rue
- American Arborvitae
- Star Jasmine
- Canada Hemlock
- Tulip species
- Veronica, Speedwell; Sunny Border
- Japanese Viburnum
- Sweet Viburnum
- Japanese Snowball

08 06 2012 Label amendment to add marketing statements and correct application rate.

- V. rigidum
- V. suspensum
- V. tinus
- V. trilobium
- V. wrightii
- Vinca major
- Vinca Minor
- Vitis spp.
- Weigela florida
- Yucca aloifolia
- Y. filametosa
- Canary Island Viburnum
- Arrowwood Viburnum
- Laurustinus
- Cranberry Bush
- Leatherleaf Viburnum
- Greater Periwinkle, Vinca
- Common Periwinkle, Vinca
- Grape*
- Old Fashioned Weigela
- Spanish Bayonet, Yucca
- Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, of, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire direction for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc election, the replacement of product.